

Idaho Falls Pediatrics

Ankle Sprain (for Teenagers) - Teen

Illustration of a sprained ankle

What is an ankle sprain?

An ankle sprain is an injury that causes a stretch or tear of one or more ligaments in the ankle joint. Ligaments are strong bands of tissue that connect bones at the joint.

Sprains may be graded I, II, or III depending on their severity:

- grade I sprain: pain with minimal damage to the ligaments
- grade II sprain: more ligament damage and mild looseness of the joint
- grade III sprain: complete tearing of the ligament and the joint is very loose or unstable.

Sometimes sprains are just classified as mild or severe, depending on the amount of ligament damage.

There are many ligaments in the ankle. The most common type of sprain involves the ligaments on the outside part of the ankle (lateral ankle sprain). Ligaments on the inside of the ankle may also be injured (medial ankle sprain) as well as ligaments that are high and in the middle of the ankle (high ankle sprains).

How does it occur?

A sprain is caused by twisting your ankle. Your foot usually turns in or under but may turn to the outside.

What are the symptoms?

Symptoms of a sprained ankle include:

- mild aching to sudden pain
- swelling
- discoloration
- inability to move the ankle properly
- pain in the ankle even when you are not putting any weight on it.

How it is treated?

Treatment may include:

- Applying ice packs to your ankle for 20 minutes. Repeat every 4 hours
- Elevating your ankle by placing a pillow underneath your foot for 1 day. Try to keep your ankle above the level of your heart.
- Wrapping an elastic bandage around your ankle to keep the swelling from getting worse. Keep the wrap on for 2 days.
- Taking anti-inflammatory pain medication such as ibuprofen.

Rarely, severe ankle sprains with complete tearing of the ligaments need surgery. After surgery your ankle will be in a cast for 4 to 8 weeks.

How long will the effects last?

The length of recovery depends on many factors such as your age, health, and if you have had a previous ankle injury. Recovery time also depends on the severity of the sprain. A mild ankle sprain may recover within a few weeks, whereas a severe ankle sprain may take 6 weeks or longer to recover. Recovery also depends on which ligaments were torn. A lateral sprain (outside ligaments) takes less time to recover than a medial sprain (inside ligaments) or a high ankle sprain (high, middle ligaments).

When can I return to my normal activities?

Everyone recovers from an injury at a different rate. Return to your activities will be determined by how soon your ankle recovers, not by how many days or weeks it has been since your injury has occurred. In general, the longer you have symptoms before you start treatment, the longer it will take to get better. The goal of rehabilitation is to return you to your normal activities as soon as is safely possible. If you return too soon you may worsen your injury.

You may safely return to your normal activities when, starting from the top of the list and progressing to the end, each of the following is true:

- You have full range of motion in the injured ankle compared to the uninjured ankle.
- You have full strength of the injured ankle compared to the uninjured ankle.
- You can walk straight ahead without pain or limping.

How can I help prevent an ankle sprain?

To help prevent an ankle sprain, follow these guidelines:

- Wear proper, well-fitting shoes when you exercise.
- Stretch gently and adequately before and after athletic or recreational activities.
- Avoid sharp turns and quick changes in direction and movement.
- Consider taping the ankle or wearing a brace for strenuous sports, especially if you have a previous injury.

*NOTE: This information is provided as a public educational service. The information does not replace any of the instructions your physician gives you. If you have a medical emergency please call 911 or call the Hospital at (208) 529-6111. If you have questions about your child's care, please call Idaho Falls Pediatrics at (208) 522-4600.