

Idaho Falls Pediatrics

Hay Fever (Allergic Rhinitis)

What is hay fever?

Hay fever is an allergic reaction of the nose (and sinuses) to pollen or another substance in the air.

Symptoms include:

- a clear nasal discharge
- an itchy nose with sneezing and sniffing
- itchy, watery eyes (eye allergies)
- sometimes, sinus or ear congestion, tickling sensation in the back of the throat.

Hay fever is the most common allergy. More than 15% of people have it.

What is the cause?

Although pollen is usually the cause of hay fever, it can also be caused by animal dander or something else your child is allergic to. This allergic sensitivity is often inherited.

During late April and May the most common pollen causing hay fever is from trees. From late May to mid-July, the pollen is usually from grass. From late August to the first frost, the leading cause of hay fever is ragweed pollen.

How long will it last?

This is a chronic condition that will probably come back every year during pollen season, perhaps for a lifetime. Therefore, it is important to learn how to control it.

How can I take care of my child?

- **Oral antihistamine medicine**

The best drug for hay fever is an antihistamine. It will relieve nose and eye symptoms.

Symptoms clear up faster if antihistamines are given at the first sign of sneezing or sniffing. For children with daily symptoms, the best control is attained if antihistamines are taken continuously throughout the pollen season. For children with occasional symptoms, antihistamines can be taken on days when symptoms are present or expected.

The main side effect of antihistamines is drowsiness. If your child becomes drowsy, switch to a combination product that contains an antihistamine with a decongestant (such as pseudoephedrine). If your child remains drowsy, continue the drug, but temporarily decrease the dosage. Your child should become tolerant of the regular dosage in 1 to 2 weeks. Newer prescription antihistamines cause much less drowsiness and are FDA approved for use in children over age 6. Examples are Allegra, Zyrtec, and Claritin.

- **Prescription steroid nasal sprays for prevention**

If not helped by antihistamines, severe hay fever can usually be controlled by prescription steroid nasal sprays. Allergy shots are usually not needed.

Nasal sprays must be used when the nose is not dripping. Give your child an antihistamine to stop the dripping before you use the spray.

- **Pollen removal to decrease symptoms of hay fever**

Pollen tends to collect on the exposed body surfaces and especially in the hair. Shower your child and wash his hair every night before he goes to bed. Your child should avoid handling pets that have been outside and are probably covered with pollen.

- **Prevention of hay fever symptoms**

Your child's exposure to pollen can be reduced by not going on drives in the country and by not sitting by an open car window on necessary drives.

He should stay away from someone cutting the grass during pollen season. When it is windy or the pollen count is especially high, he should stay indoors. Close the windows that face the prevailing winds. Use an air conditioner rather than an attic or window fan. Fans can pull in pollen.

If your child's hay fever is especially bad, you could also take him to an air-conditioned store or theater for a few hours.

Avoid feather pillows, pets, farms, stables, and tobacco smoke if any of them seem to bring on symptoms of nasal allergy.

If your child is allergic to ragweed, you may wish to plan a vacation to an area that has little or no ragweed. Only the coastal areas of Washington, Oregon, and California are free of ragweed pollen.

- **Eye allergies associated with hay fever**

If your child also has itchy, watery eyes, wash his face and eyelids to remove pollen. Then apply a cold wet cloth to the eyelids for 10 minutes. An oral antihistamine will usually bring the eye symptoms under control. If not, put 1 drop of long-acting decongestant eyedrops (a nonprescription item) in the eyes every 8 to 12 hours for a few days. Ask your pharmacist to recommend a reliable product.

- **Common mistakes**

Decongestant nosedrops or nasal sprays usually do not help hay fever because they are washed out by nasal secretions as soon as they are put into the nose. Also, if decongestant nosedrops or nasal sprays are used for more than 5 days, they can irritate the nose and make it more congested.

When should I call Idaho Falls Pediatrics?

Call during office hours if:

- Your child's symptoms are not controlled in 2 days with antihistamines.
- Your child develops sinus pain or pressure.
- You have other concerns or questions.

*NOTE: This information is provided as a public educational service. The information does not replace any of the instructions your physician gives you. If you have a medical emergency please call 911 or call the Hospital at (208) 529-6111. If you have questions about your child's care, please call Idaho Falls Pediatrics at (208) 522-4600.